

<b>Products</b>	Biovac® direct suction device
<b>Procedural Area</b>	Bleed Management
<b>Article</b>	Immediate unprepared hydroflush colonoscopy for severe lower GI bleeding: a feasibility study
<b>Publication</b>	Gastrointestinal Endoscopy 2012 Volume 76, No. 2
<b>URL</b>	<a href="http://dx.doi.org/10.1016/j.gie.2012.03.1391">http://dx.doi.org/10.1016/j.gie.2012.03.1391</a>
<b>Author</b>	Aparna Repaka, MD, Matthew R. Atkinson, MD, Ashley L. Faulx, MD, Gerard A. Isenberg, MD, MBA, Gregory S. Cooper, MD, Amitabh Chak, MD, Richard C.K. Wong, MBBS
<b>Purpose</b>	To evaluate the practicality, safety, and results of using hydroflush colonoscopy to prepare the colon during an emergency lower GI bleed
<b>Key Points</b>	<p><u>BACKGROUND</u></p> <ul style="list-style-type: none"> <li>- Urgent colonoscopy requires an immediate oral bowel preparation, which delays diagnosis and therapy</li> <li>- The use of water jets and mechanical suction devices can potentially improve visualization in a timelier manner</li> </ul> <p><u>STUDY METHODS</u></p> <ul style="list-style-type: none"> <li>- Hydroflush colonoscopy (combination of water-jet pump irrigation and the <b>Biovac® direct suction device</b>) was used to cleanse the colon</li> <li>- Performed during 13 procedures on patients with bloody bowel movement within the past 24 hours, admission to the intensive care unit, and other criteria</li> <li>- Outcomes were measured primarily by the percentage of hydroflush procedures that resulted in a satisfactory examination of the entire colon</li> <li>- Visualization of the bleeding site, ICU stay, hospital stay, recurrent bleed rates and transfusion requirements were also measured</li> </ul> <p><u>RESULTS</u></p> <ul style="list-style-type: none"> <li>- <b>100% of the procedures were deemed to have adequate visualization</b></li> <li>- 69.2% of the procedures resulted in a complete examination of the colon</li> <li>- A diagnosis (presumptive/definite) was made in 100% of the procedures</li> <li>- A definitive diagnosis was made in 38.5% of the procedures</li> <li>- Median ICU and hospital stay were 1.5 and 4.3 days, respectively</li> <li>- 25% of patients experienced recurrent bleeding during hospital stay</li> </ul>
<b>Conclusions</b>	Hydroflush colonoscopy in cases of severe lower GI bleeding is practical and may reduce time to endoscopy, increase diagnostic outcomes and rate of therapy
<b>This page does not replace a subscription.</b>	